

# Resistance training in musculoskeletal rehabilitation: a systematic review

Jakob Kristensen, Andy Franklyn-Miller

Centre for Human Performance, Rehabilitation and Sports Medicine, Defence Medical Rehabilitation Centre Headley Court, Epsom, Surrey, UK

## Correspondence to

Jakob Kristensen, Centre for Human Performance, Rehabilitation and Sports Medicine, Defence Medical Rehabilitation Centre Headley Court, Epsom, Surrey KT18 6JN, UK; jakob.kristensen711@mod.uk

Accepted 16 June 2011

Published Online First

26 July 2011

## ABSTRACT

**Objective** To review the efficacy of resistance training (RT) as a therapeutic modality in various musculoskeletal conditions.

**Design** Systematic review.

**Subjects** Data from 1545 rehabilitation patients who had participated in structured RT programmes were included into the review. The total number of patients was composed of separate musculoskeletal conditions—chronic low back pain (CLBP) (549), tendinopathy (299), knee osteoarthritis (433), anterior cruciate ligament reconstruction (189) and hip replacement surgery (75).

**Results** Evidence suggests that RT can increase muscle strength, reduce pain and improve functional ability in patients suffering from CLBP, knee osteoarthritis, and chronic tendinopathy and those under recovery after hip replacement surgery.

**Conclusion** RT can be used successfully as a therapeutic modality in several musculoskeletal conditions, especially those of a chronic variety. Although the exact application of training intensity and volume for maximal therapeutic effects is still unclear, it appears that RT guidelines, which have proven effective in a healthy population, can also be successfully applied in a rehabilitation context.

## INTRODUCTION

The effects of musculoskeletal resistance training (RT) on the development of strength and power in a healthy population have been well covered in the literature.<sup>1–4</sup> Specifically, RT can result in increased muscle size, maximal strength and muscle power through several mechanisms. These range from skeletal muscle hypertrophy and changes in muscle architecture to neural adaptations such as increased motor unit activation and supraspinal adaptations.<sup>2 5–8</sup> However, relatively little is known about the effects of RT in an injured population and, moreover, even less is known about what constitutes the optimal guidelines for its use. In recent years, the increased prevalence of RT in various rehabilitation programmes lends some empirical evidence to its beneficial effects in a rehabilitation context; however, the scientific evidence for the use of RT in rehabilitation is not as extensive as the evidence presented in a healthy population. The objective of this review was to summarise the effects of RT in a rehabilitation context with regards to its effects on maximal strength, functional ability, alleviation of pain and quality of life (QoL) parameters in the musculoskeletal conditions, where RT is most commonly prescribed throughout the literature. Furthermore,

this review will also seek to critically evaluate the validity of the RT methods most commonly used in musculoskeletal rehabilitation programmes.

## METHODS

A comprehensive search of the PUBMED/MEDLINE, CINAHL and SportDiscus databases was conducted by the authors. All publications in the English language listed up until April 2010 (including online format only) were considered for inclusion. Abstracts were initially screened and then full-text versions accessed when included.

The following inclusion criteria were adhered to: (1) studies with an intervention period lasting minimum 4 weeks, (2) studies with more than one clinically relevant outcome measure, (3) studies using patients suffering from a clinically diagnosed musculoskeletal condition and (4) studies using external resistance in addition to bodyweight as part of the RT intervention. Both randomised controlled trials and observational studies were included into the review.

The comprehensive database search based on relevant MeSH terms (RT, chronic low back pain (CLBP), knee osteoarthritis, achilles/patellar tendinopathy, anterior cruciate ligament (ACL) reconstruction, hip replacement surgery) identified a number of studies (figure 1).


A summary of the changes in outcome measures in the papers included into this review can be found in table 1.

## RESULTS

### Chronic low back pain

Chronic low back pain (CLBP) is the most common musculoskeletal condition affecting a wide array of people from both athletic and non-athletic backgrounds. In most cases, the aetiology of the condition appears to be multifactorial but, nevertheless, is associated with a deconditioning of the extensor muscles of the back and resultant loss of muscle strength,<sup>9–12</sup> thus creating a potential target for RT in CLBP rehabilitation.

RT has been shown to improve strength<sup>13–17</sup> and reduce self-reported pain in patients with CLBP,<sup>14 17</sup> thereby serving as an effective therapeutic modality in this common condition.<sup>18 19</sup> Long-term benefits after RT intervention can be observed in patients with CLBP<sup>20</sup> but also appear to be further influenced by extrinsic factors such as physical activity, smoking and treatment outcome expectations.<sup>21</sup> Isolated data have suggested that RT offers no benefits over normal physical therapy in the treatment of CLBP;<sup>22</sup> however,



**Figure 1** Search strategy. Numbers of studies either used or rejected are indicated in brackets.

such conclusions are not supported by the majority of the current literature.

The details of what constitutes an optimal RT programme in CLBP such as training intensity, volume and contraction type are largely unknown at present, as evidenced by the widely differing RT methodology employed by different studies.<sup>13–18</sup> Sertpoyraz *et al*<sup>23</sup> saw increased lumbar extension strength and reduced self-reported pain in patients after participation in an isokinetic RT programme; however, the results were not significantly different when compared with a standard therapeutic exercise programme. Unfortunately, the authors failed to provide a description of the standard exercise programme, thus making a direct comparison of the two training programmes difficult.

With regards to RT volume in the treatment of CLBP, Limke *et al*<sup>24</sup> reported that completing two sets rather than one set of a RT exercise did not lead to increased benefits either on measures of strength or self-reported pain. This is directly contradictory to evidence from a healthy population where meta-analysis has confirmed the superiority of a multiple-set approach for eliciting maximal strength gains.<sup>25–28</sup> However, the study by Limke *et al*<sup>24</sup> was strictly speaking not a single-set study design as the authors did make use of five to six different strengthening exercises in their training protocol leading to a higher accumulated training volume.

With regards to RT intensity, Helmhout *et al*<sup>20</sup> compared high- and low-intensity lumbar strengthening programmes showing greater strength gains with a high-intensity approach but less reduction in kinesiophobia, prompting the authors to conclude that high-intensity RT offered no added benefits over low-intensity RT in restoring back function in CLBP. Similarly, Harts *et al*<sup>17</sup> found that participation in a high-intensity RT programme did not lead to greater strength gains when compared with a low-intensity programme; however, it did seem to improve patient-reported QoL more. However, in both these above studies, the definition of high-intensity RT was different to the conventional definition of high-intensity RT in a healthy population. It is widely accepted that an exercise intensity >70% of 1RM is the threshold above which neuromuscular adaptations occur in response to RT and where the training can be classified as high-intensity.<sup>29–32</sup> The methodology of Harts *et al*<sup>17</sup> consisted of a ‘high-intensity’ group exercising at 50% of 1RM, an exercise intensity which is insufficient for eliciting strength gains in a healthy population.

Unsurprisingly, the authors failed to see significant strength gains even in the ‘high-intensity’ group, indicating that sufficient exercise intensity is needed to elicit neuromuscular adaptations even in an injured population. Incidentally, the study by Helmhout *et al*,<sup>22</sup> which concluded that RT offered no benefits over normal physical therapy in the treatment of CLBP, used the same insufficient exercise intensity in the RT protocol.

Recently, periodised RT, which is the most effective form of RT in a healthy population,<sup>32</sup> has also been shown to be effective at increasing muscular strength, reducing pain and improving QoL in patients with CLBP.<sup>33–34</sup> In a study by Kell *et al*,<sup>33</sup> CLBP patients engaged in either a 16-week whole-body periodised RT programme using an exercise intensity range of 53–72% of 1RM or a periodised aerobic training programme. Interestingly, only the RT programme resulted in a reduction of pain scores and improvements in QoL parameters. In a follow-up study by Jackson *et al*,<sup>34</sup> CLBP patients used an exercise intensity ranging from 55% to 79% of 1RM. Again, increased muscle strength, reduced pain and increased functional ability were evident after participation in the RT programme, which was well-tolerated by all patients.<sup>34</sup> These results indicate that improving maximal muscle strength through an effective RT programme can reduce symptoms in CLBP patients and that such an approach may be considered a valid therapeutic modality.

### Chronic tendinopathy

Chronic tendinopathy is prevalent in both recreational and elite athletes, especially in those who engage in sporting activities with a heavy emphasis on running and jumping. In recent years, the potential use of RT as a therapeutic modality in the treatment of chronic tendinopathy has received increased interest. The majority of this interest has focused on the use of eccentric-only exercise to reduce pain and improve QoL of those affected by the condition. This type of RT uses only the eccentric or ‘muscle-lengthening’ component of muscle contractions to load the muscles being exercised.

Particularly in chronic Achilles tendinopathy, the results seem promising. In this condition, eccentric-only RT has been associated with decreased pain and improved function immediately after exercise intervention.<sup>35–39</sup> In one study, these improvements were still observed at long-term follow-up at

**Table 1** Papers included into the systematic review

Study	Number of participants (n)	Mean age (years)	Change in outcome measures
<b>Chronic low back pain</b>			
Manniche, 1991	35	45	↓PS
Risch <i>et al</i> , 1993	31	45	↑MS, ↓PS, ↑FA
Rissanen <i>et al</i> , 1995	30	40	↑MS, ↑CSA
Danneels <i>et al</i> , 2001	40	44	↑MS, ↑CSA
Harts <i>et al</i> , 2008	20	44	→MS, ↑QoL
Helmhout <i>et al</i> , 2004	40	41	↑FA, ↑MS, ↑QoL
Petersen <i>et al</i> , 2007	130	35	↑FA, ↓PS
Helmhout <i>et al</i> , 2008	64	37	↑FA, ↓PS
Limke <i>et al</i> , 2008	100	47	↑MS, ↓PS, ↑FA
Sertpoyraz <i>et al</i> , 2009	20	39	↓PS, ↑MS, ↑FA
Kell <i>et al</i> , 2009	9	40	↑MS, ↓PS, ↑QoL, ↑BC
Jackson <i>et al</i> , 2010	30	52 <sup>1</sup> , 63 <sup>2</sup>	↑MS, ↓PS, ↑FA, ↑QoL
<b>Tendinopathy</b>			
Alfredson <i>et al</i> , 1998	15	44	↑MS, ↓PS
Mafi <i>et al</i> , 2001	44	48	↓PS
Roos <i>et al</i> , 2004	15	45	↑FA, ↓PS
Langberg <i>et al</i> , 2007	12	26	↓PS
Mafulli <i>et al</i> , 2008	45	26 <sup>1</sup> , 28 <sup>2</sup>	↓PS, ↑FA
Gardin <i>et al</i> , 2010	20	49	↑FA, ↓PS
Silbernagel <i>et al</i> , 2001	22	45	↓PS, ↑FA
Purdam <i>et al</i> , 2004	9	28	↓PS
Young <i>et al</i> , 2005	17	27	↑FA, ↓PS
Frohm <i>et al</i> , 2007	20	26 <sup>1</sup> , 28 <sup>2</sup>	↓PS, ↑FA, ↑MS
Visnes <i>et al</i> , 2005	13	27	FA
Jonsson and Alfredson, 2005	15	25	↓PS, ↑FA
Bahr <i>et al</i> , 2006	10	31	↑MS, ↑FA
Kongsgaard <i>et al</i> , 2009	26	31 <sup>1</sup> , 32 <sup>2</sup>	↑MS, ↓PS, ↑FA
Kongsgaard <i>et al</i> , 2010	8	33	↓PS, ↑FA
<b>Knee osteoarthritis</b>			
Ettinger <i>et al</i> , 1997	147	68	↑FA, ↓PS, ↑MS
Gur <i>et al</i> , 2002	17	55 <sup>1</sup> , 56 <sup>2</sup>	↑MS, ↑FA, ↓PS, ↑CSA
Topp <i>et al</i> , 2002	67	66 <sup>1</sup> , 64 <sup>2</sup>	↑FA, ↓PS
Schilke <i>et al</i> , 1996	10	65	↑MS, ↑FA, ↓PS
Mikesky <i>et al</i> , 2006	113	69	(↓)MS
King <i>et al</i> , 2008	14	48	↑MS, ↑FA
Jan <i>et al</i> , 2008	68	63 <sup>1</sup> , 62 <sup>2</sup>	↑FA, ↓PS, ↑MS
<b>Anterior cruciate ligament reconstruction</b>			
Mikkelsen <i>et al</i> , 2000	44	26 <sup>1</sup> , 19 <sup>2</sup> , 25 <sup>3</sup> , 19 <sup>4</sup>	↑MS
Morrissey <i>et al</i> , 2002	43	29 <sup>1</sup> , 28 <sup>2</sup>	↓PS, ↑MS
Holm <i>et al</i> , 2006	26	26	↑MS, ↑CSA
Liu-Ambrose <i>et al</i> , 2003	5	25	↑MS, ↑FA
Risberg <i>et al</i> , 2007	35	27 <sup>1</sup> , 31 <sup>2</sup>	↓PS, (↓)MS
Gerber <i>et al</i> , 2007 <sup>83</sup>	16	29 <sup>1</sup> , 31 <sup>2</sup>	↑MS, ↑FA
Gerber <i>et al</i> , 2007 <sup>84</sup>	20	29	↑CSA
<b>Hip replacement</b>			
Hauer <i>et al</i> , 2002	15	82	↑MS, ↑FA
Suetta <i>et al</i> , 2004 <sup>90</sup>	13	69	↑MS, ↑FA, ↑CSA
Suetta <i>et al</i> , 2004 <sup>91</sup>	12	71	↑MS, ↑CSA
Suetta <i>et al</i> , 2008	12	71	↑MS, ↑CSA, ↑FA
Husby <i>et al</i> , 2009	12	58	↑MS, ↑FA, ↑QoL
Husby <i>et al</i> , 2010	11	58	↑MS, ↑FA

Arrows indicate changes in outcome measures. Numbers in superscript indicate separate intervention groups. BC, body composition; CSA, cross-sectional area; FA, functional ability; MS, muscle strength; PS, pain score; QoL, quality-of-life parameters.

1-year postintervention,<sup>37</sup> and in another study continued improvements were seen more than 4-year postintervention.<sup>40</sup> Some evidence suggest that eccentric-only training may offer additional pain-relieving benefits over concentric-only

training,<sup>36</sup> and eccentric-concentric training.<sup>41</sup> However, the methodology of several of these eccentric-only studies has been criticised, leading some authors to conclude that the possible beneficial effects of eccentric loading as a

therapeutic modality in Achilles tendinopathy are still not fully documented.<sup>42–43</sup>

Currently, it is unclear what constitutes an effective eccentric training volume for eliciting optimal effects,<sup>44</sup> and doubt has also been cast on whether these eccentric-only training protocols can be classified as RT per se. Although the majority of studies have used external resistance in addition to bodyweight as part of the training progression, the exercise intensity used in these studies is generally below the level considered necessary for eliciting strength gains through neural and morphological adaptations, at least in a healthy population.<sup>1–29–31</sup> In fact, it has recently been proposed that the beneficial effects of eccentric-only exercise in Achilles tendinopathy are the result of stretching of passive structures rather than actual muscle strengthening through eccentric loading.<sup>45</sup> Currently, no research has investigated if high-intensity RT, utilising eccentric/concentric muscle actions has therapeutic benefits in the treatment of chronic Achilles tendinopathy.

RT has also been investigated as a potential therapeutic modality in patients suffering from patellar tendinopathy. However, here the literature is less extensive and more inconsistent. Some studies have reported improved function and pain reduction after eccentric-only exercise.<sup>46–48</sup> On the contrary, Visnes *et al*<sup>49</sup> reported no pain-reducing effects of eccentric-only exercise in competitive athletes suffering from patellar tendinopathy, at least when performed in the competitive season. As in Achilles tendinopathy, only very limited evidence exists to suggest that eccentric-only quadriceps training is more effective than other types of RT at alleviating pain in patients suffering from patellar tendinopathy.<sup>50</sup> However, when compared with more aggressive treatments such as surgery, eccentric-only exercise appeared no better at reducing patient-reported pain and only ~35% of patients were symptom-free 1 year after the exercise intervention.<sup>51</sup> Taken together, these results indicate that the use of eccentric-only training in the management of patellar tendinopathy is associated with a degree of uncertainty with regards to treatment outcome.

Very little data exist on the effects of conventional eccentric-concentric RT in the treatment of patellar tendinopathy. A recent study by Kongsgaard *et al*<sup>52</sup> reported superior effects after 12 weeks of heavy slow RT when compared with eccentric-only training in patients suffering from patellar tendinopathy. Both groups saw improvements in patient-reported pain; however, only heavy slow RT led to improvements in tendon pathology. These positive effects included reductions in tendon swelling and vascularisation along with increased collagen turnover. These findings are interesting because this is the only study in patellar tendinopathy which has used an exercise intensity of sufficient magnitude to elicit neural and morphological adaptations.<sup>29–31</sup> The above-mentioned study used a variety of RT exercises to specifically target and strengthen the muscles around the knee and hip joint, and, interestingly, this general strengthening approach did appear to effectively alleviate the symptoms of patellar tendinopathy. Furthermore, at the 6-month follow-up, the patients who had completed the heavy RT programme were more satisfied with their treatment outcome than those who had undergone an eccentric-only training programme. In a recent follow-up study, Kongsgaard *et al*<sup>53</sup> reported that heavy slow RT improved abnormal tendon morphology in patellar tendinopathy, leading the authors to hypothesise that this is the potential mechanism of action by which heavy RT improves the clinical outcome of this chronic condition. Thus, these

two recent studies indicate that participation in a heavy RT programme may offer additional benefits over both eccentric-only and low-intensity RT in patients suffering from patellar tendinopathy by improving a wider range of outcome measures. Also, based on the current literature in chronic lower limb tendinopathy, it appears that RT is effective at reducing symptoms in both young and old patients alike and that this effectiveness persists regardless of gender.

### Knee osteoarthritis

Knee osteoarthritis is a degenerative condition which predominantly affects the middle-aged and older population. However, other key risk factors include obesity and prior sports-related knee injury. Weakness of the quadriceps muscle has been shown to correlate significantly with both functional ability and pain in knee osteoarthritis.<sup>54–55</sup> Thus, strengthening of the quadriceps muscle with RT potentially offers benefits in the treatment of knee osteoarthritis.

It has been reported that patients who participate in RT show improvements in pain and functional ability.<sup>56–60</sup> Participation in a RT programme can attenuate the progressive loss of muscle strength commonly seen in older knee osteoarthritis patients and may retard disease progression.<sup>61</sup> Some evidence suggests that the positive therapeutic effects of RT are evident even in patients with advanced knee osteoarthritis.<sup>62</sup> However, these studies have used a variety of different RT methods, and it appears unclear what constitutes the optimal training methods to employ in the treatment of knee osteoarthritis.<sup>59</sup>

Gur *et al*<sup>57</sup> used isokinetic RT, which due to the fixed angular velocities does not mimic the contraction pattern of real-life movements, and is generally viewed as less functional than dynamic isotonic RT.<sup>29–31</sup> Despite this notion, the authors still saw significant improvements in muscular strength and functional ability with isokinetic RT. However, the authors<sup>57</sup> did observe better results from combined eccentric-concentric training than from concentric-only RT, indicating that methods which use the conventional contraction coupling, that is, the stretch-shortening cycle, may be more effective in the treatment of knee osteoarthritis. Interestingly, the latter seems to be challenged by Topp *et al*,<sup>58</sup> who saw no increased therapeutic benefit of dynamic over isometric resistance exercise. In this particular study, both types of RT resulted in similar improvements in patient-reported pain and functional ability. The improvements in muscular strength from isometric training are generally thought to be angle specific;<sup>29–31</sup> however, Topp *et al*<sup>58</sup> used isometric training at a variety of joint angles, thereby strengthening the musculature in a method akin to a full range of motion. This potentially explains the beneficial effects of isometric RT in the study.

Furthermore, it appears unclear what exercise intensity is needed to observe optimal results from RT when used as a therapeutic modality in knee osteoarthritis. Although not significant, Jan *et al*<sup>63</sup> observed a trend towards better results with high-intensity RT with regards to improving strength, reducing pain and improving functional ability. In this particular study, the high-intensity group exercised at an intensity of 60% of 1RM, which although lower than what is recommended for optimal results in a healthy population,<sup>32</sup> is potentially still sufficient to elicit some neuromuscular adaptations.<sup>29–31</sup> The authors suggested that resistance exercise of >80% of 1RM is not feasible in knee osteoarthritis patients due to pain and possible detrimental effects associated with repetitive heavy loading, a view that is supported by others in theory.<sup>64</sup> However,

other evidence suggests that heavy RT does not exacerbate joint problems in middle-aged and older individuals,<sup>5</sup> and Andersen *et al*<sup>65</sup> even speculated that the inclusion of heavy RT exercises in a knee rehabilitation programme would lead to superior results owing to the increased neuromuscular activation observed when compared with conventional therapeutic rehabilitation exercises. Lange *et al*<sup>66</sup> reported that participation in a progressive high-intensity RT programme, using an exercise intensity of ~80% of 1RM retarded disease progression in knee osteoarthritis patients by favourably impacting cartilage morphology, thereby disproving the notion that high-intensity RT is not feasible in knee osteoarthritis rehabilitation.

Although some evidence suggests that RT above 60% of 1RM may offer beneficial therapeutic effects in the treatment of knee osteoarthritis, additional research is needed to conclusively establish if this method offers additional benefits over lighter, less-intense RT in the treatment of knee osteoarthritis.

### Rehabilitation after ACL reconstruction

RT has been studied comprehensively as a potential adjunct therapy after knee surgery. Following ACL reconstruction surgery, RT is routinely prescribed as part of the rehabilitation process.<sup>67–69</sup> Loss of quadriceps muscle strength and joint range of motion are two well-established complications of ACL reconstruction surgery,<sup>70–73</sup> and this weakness can persist for years after surgery if not adequately addressed by an effective rehabilitation programme.<sup>70 71 74</sup> Consequently, some authors emphasise that restoring quadriceps strength is vital to a successful therapeutic outcome,<sup>68 75 76</sup> and some evidence suggests that RT is effective at improving strength, functional ability and reducing pain in patients postsurgery,<sup>77–79</sup> however, not more so than a proprioception training programme<sup>80</sup> or neuromuscular training programme.<sup>81</sup> Interestingly, other types of exercise such as stair climbing and cycle ergometry also appear to lead to significant gains in quadriceps muscle strength when used in postoperative rehabilitation following ACL reconstructive surgery.<sup>82</sup> This indicates that postsurgery, RT offers no additional benefits over other conventional types of exercise in promoting a return of muscle strength and functional ability, at least in the short term.

The use of eccentric-only resistance exercise has also been investigated as potential treatment modality after ACL reconstruction. Gerber *et al*<sup>83 84</sup> reported that eccentric-only RT resulted in greater strength gains, daily activity level and quadriceps muscle hypertrophy when compared with a standard rehabilitation protocol, with the beneficial effects persisting 1 year postsurgery.<sup>85</sup> However, in the above studies, the eccentric RT consisted of eccentric cycle ergometry with no definitive description of the intensity level used, making it difficult to actually classify this intervention as an actual RT programme. Thus, based on the above studies, it is not possible to conclude whether eccentric-only RT offers additional benefits over traditional RT methods in the rehabilitation period following ACL reconstruction surgery.

As even low-to-moderate-intensity exercise results in significant strength gains after ACL reconstruction surgery, there is little to be gained from the use of high-intensity RT in the immediate period following ACL reconstruction surgery. However, it is not clear whether this type of RT could potentially benefit patients in the long-term, especially considering that persistent muscle weakness is evident in many patients, sometimes even years after participation in traditional rehabilitation programmes.<sup>69 70 73</sup> Therefore, additional research

is needed to investigate if high-intensity RT can be introduced successfully into a rehabilitation protocol at a later stage and potentially improve the long-term prognosis for patients who have undergone ACL reconstruction surgery.

### Rehabilitation after hip replacement surgery

Hip osteoarthritis is a degenerative condition, which particularly affects the older population and often leads to the need for hip replacement surgery. Recent data suggest that this procedure is being carried out with increasing frequency in the UK,<sup>86</sup> thus creating a need for increasingly efficient rehabilitation strategies. Evidence suggests that hip replacement patients often fail to fully regain the muscle strength and functional ability lost due to the hospitalisation/immobilisation process.<sup>87 88</sup> Consequently, it has been speculated that RT, due to its ability to increase maximal muscle strength, may ameliorate this postsurgery strength loss and thus serve as a distinct therapeutic modality.

Participation in an RT programme increases maximal muscle strength in patients having undergone hip replacement surgery.<sup>89–94</sup> This increase in maximal strength has been observed as early as 4–5 weeks after the start of an RT programme,<sup>90 91 93</sup> with some of the strength gains and neuromuscular adaptations still evident 11 months after cessation of the RT programme.<sup>93</sup> Similarly, a range of functional performance parameters such as walking speed, stair climbing and seated-to-standing time also appear to be significantly improved by RT after hip replacement surgery.<sup>89 90</sup> Suetta *et al*<sup>90</sup> compared the effects of RT, percutaneous neuromuscular electrical stimulation (NMES) and conventional rehabilitation in patients having undergone hip replacement surgery. When all three training methods were commenced while patients were still in hospital recovering after surgery, only RT resulted in a significantly reduced length of stay in hospital. Similarly, only RT prevented postsurgery muscle atrophy and was the only intervention resulting in increased muscle cross-sectional area and maximal muscle strength 12 weeks postsurgery. However, it is worth noting that both the RT and NMES groups showed significant improvements in functional performance tests such as walking speed and stair-climbing 12 weeks postsurgery, indicating some merit to the use of NMES in the postsurgery recovery period. In the study by Suetta *et al*,<sup>90</sup> the RT protocol consisted of a progressive increase in training intensity from 65% of 1RM immediately after hospital discharge up to 80% of 1RM for the last 6 weeks. Thus, the training intensity eventually surpassed the theoretical minimum for neuromuscular adaptations to take place, and, not surprisingly, the last 6 weeks of the study also saw the greatest increases in peak force.

Another study by Suetta *et al*<sup>91</sup> found that RT was the only training method which resulted in significant increases in muscle rate of force development (RFD). This RT-induced increase in RFD has been corroborated by recent studies<sup>93 94</sup> and is potentially linked to evidence suggesting that older hip replacement patients experience proportionately greater increases in fast-twitch type IIx muscle fibre area in response to RT.<sup>92</sup> The increase in RFD seems particularly relevant from a functional performance point of view and has led some to conclude that RT would make older patients more able to prevent falls due to their increased ability to rapidly generate muscle force in situations where balance is lost.

When used as a therapeutic modality after hip replacement surgery, it is evident that high-intensity RT (>70% of 1RM) can be used successfully to elicit improvements in

maximal strength, RFD, muscle morphology and functional performance.<sup>89–94</sup> It appears that these beneficial results can be achieved even in older patients (>60 years) who take longer to regain strength after a period of disuse/immobilisation than younger individuals.<sup>95</sup> Nevertheless, the positive effects of RT are still evident despite an advanced age.

The results are evident either by utilising a progressive increase in training intensity before attempting heavier loads,<sup>90–92</sup> or by using high-intensity RT immediately postdischarge without a progressive build-up.<sup>89 93 94</sup> Both approaches appear to be equally well tolerated by hip replacement patients, although there is some evidence to suggest that the sooner the patients start to train using a high intensity, the quicker the increase in maximal strength is manifested.<sup>89 93</sup> Taken together, these results indicate that the use of RT immediately postoperatively in hip replacement patients shortens the time before hospital discharge and that continuing RT post-discharge can increase maximal strength, RFD and functional performance even in an advanced age group.

## DISCUSSION

Based on existing literature, RT is a useful tool in the rehabilitation of a variety of musculoskeletal conditions, especially those where loss of muscular strength and functional ability is evident (see table 1). As such, the beneficial effects of RT are apparent in chronic conditions such as a recurrent low back pain, knee osteoarthritis and patellar tendinopathy, whereas the effectiveness of RT in a postsurgical setting seems to vary. High-intensity RT does not appear feasible post ACL reconstruction surgery to avoid stressing the knee joint and consequently jeopardising the integrity of the new graft. However, high-intensity RT shows clear beneficial effects when commenced very shortly after hip replacement surgery. Part of the explanation for this is potentially that after hip replacement surgery, the high-intensity RT protocols found in the existing literature predominantly stress the muscles of the knee joint (quadriceps) and exercises are performed in a supine position to avoid hip luxation. Thus, the success of RT in increasing muscle strength and functional ability after hip replacement surgery is potentially attributable to the fact that physical stress on the new hip joint is kept to a minimum, while the surrounding muscles are strengthened through contractile activity. Future work need to determine whether such a paradigm could make RT equally effective after ACL reconstruction, that is, by performing exercises which predominantly stress the hip rather than the knee joint.

Reduced capacity to adapt to a given exercise stimulus is a complication of advanced age; however, the positive effects of RT in a rehabilitation context are evident even in patients of advanced age. Although it is likely that younger patients still respond more favourably to RT than their older counterparts, RT can still be used across all ages as an effective therapeutic tool.

The current notion that RT in a rehabilitation context should be less intense than what is successfully used in a healthy population is based on the concern that high-intensity RT is potentially injurious and detrimental in an already injured population. Given the therapeutic context this is a legitimate concern, however, the recent studies which have used high-intensity RT protocols (~70% of 1RM) in the rehabilitation of musculoskeletal injuries have shown that this approach is well tolerated by patients and clearly ameliorates rather than exacerbates symptoms.<sup>33 34 66 89–94</sup> However, a key point in the safe application of high-intensity RT in the rehabilitation of musculoskeletal injuries may be the use of periodised RT. Incidentally,

## What is already known on this topic

It is already known that resistance training (RT) is a useful tool in the rehabilitation of musculoskeletal injuries and that its use may lead to a more successful treatment outcome.

## What this study adds

This study reviews the current literature regarding the effectiveness of RT in the treatment of some of the most common musculoskeletal conditions. In addition it compares which RT variables (intensity, volume etc.) are important for a successful training outcome in rehabilitation.

this approach to RT is also the most effective form of resistance exercise in a healthy population.<sup>30 32 96 97</sup> Periodised RT in rehabilitation will allow patients to gradually improve their maximal strength and become accustomed to handling heavier loads. As previously discussed, this approach has been shown to be effective at improving muscular strength and functional ability and reducing patient symptoms.<sup>33 34 90–93</sup> Although more research is needed to further elucidate what constitutes optimal RT protocols in musculoskeletal rehabilitation, new findings indicate that the principles of effective RT programme design, which have originated in a healthy population, can also be applied successfully in an injured population, despite the inherent concerns by the majority of authors.

## MAJOR FINDINGS

- ▶ RT is a valid therapeutic tool in the treatment of the most common musculoskeletal injuries, especially those of a chronic variety.
- ▶ RT is effective across age and gender.
- ▶ A high-intensity approach (>70% of 1RM) appears to be more effective than a low-intensity approach.
- ▶ High-intensity RT training does not increase the likelihood of injury, provided that patients are gradually introduced to heavier loads through periodised RT.

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

1. **McDonagh MJ**, Davies CT. Adaptive response of mammalian skeletal muscle to exercise with high loads. *Eur J Appl Physiol Occup Physiol* 1984;**52**:139–55.
2. **Aagaard P**, Simonsen EB, Andersen JL, *et al*. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *J Appl Physiol* 2002;**93**:1318–26.
3. **Aagaard P**, Thorstensen A. Neuromuscular aspects of exercise: adaptive responses evoked by strength training. In: Kjaer M, Krogsgaard M, Magnusson P, Engelbrechtsen L, Roos H, Takala T, Woo SLY, eds. *Textbook of Sports Medicine*. London: Blackwell 2003:70–106.
4. **Folland JP**, Williams AG. The adaptations to strength training: morphological and neurological contributions to increased strength. *Sports Med* 2007;**37**:145–68.
5. **Häkkinen K**, Häkkinen A. Neuromuscular adaptations during intensive strength training in middle-aged and elderly males and females. *Electromyogr Clin Neurophysiol* 1995;**35**:137–47.
6. **Aagaard P**, Andersen JL, Dyhre-Poulsen P, *et al*. A mechanism for increased contractile strength of human pennate muscle in response to strength training: changes in muscle architecture. *J Physiol (Lond)* 2001;**534**:613–23.
7. **Aagaard P**, Simonsen EB, Andersen JL, *et al*. Neural adaptation to resistance training: changes in evoked V-wave and H-reflex responses. *J Appl Physiol* 2002;**92**:2309–18.

8. **Falvo MJ**, Sirevaag EJ, Rohrbaugh JW, *et al.* Resistance training induces supraspinal adaptations: evidence from movement-related cortical potentials. *Eur J Appl Physiol* 2010;**109**:923–33.
9. **Hultman G**, Nordin M, Saraaste H, *et al.* Body composition, endurance, strength, cross-sectional area, and density of MM erector spinae in men with and without low back pain. *J Spinal Disord* 1993;**6**:114–23.
10. **Mooney V**, Gulick J, Perlman M, *et al.* Relationships between myoelectric activity, strength, and MRI of lumbar extensor muscles in back pain patients and normal subjects. *J Spinal Disord* 1997;**10**:348–56.
11. **Dvir Z**, Keating JL. Trunk extension effort in patients with chronic low back dysfunction. *Spine* 2003;**28**:685–92.
12. **Smeets RJ**, Wade D, Hidding A, *et al.* The association of physical deconditioning and chronic low back pain: a hypothesis-oriented systematic review. *Disabil Rehabil* 2006;**28**:673–93.
13. **Manniche C**, Lundberg E, Christensen I, *et al.* Intensive dynamic back exercises for chronic low back pain: a clinical trial. *Pain* 1991;**47**:53–63.
14. **Risch SV**, Norvell NK, Pollock ML, *et al.* Lumbar strengthening in chronic low back pain patients. Physiologic and psychological benefits. *Spine* 1993;**18**:232–8.
15. **Rissanen A**, Kalimo H, Alaranta H. Effect of intensive training on the isokinetic strength and structure of lumbar muscles in patients with chronic low back pain. *Spine* 1995;**20**:333–40.
16. **Danneels LA**, Vanderstraeten GG, Cambier DC, *et al.* Effects of three different training modalities on the cross sectional area of the lumbar multifidus muscle in patients with chronic low back pain. *Br J Sports Med* 2001;**35**:186–91.
17. **Harts CC**, Helmhout PH, de Bie RA, *et al.* A high-intensity lumbar extensor strengthening program is little better than a low-intensity program or a waiting list control group for chronic low back pain: a randomised clinical trial. *Aust J Physiother* 2008;**54**:23–31.
18. **Manniche C**. Clinical benefit of intensive dynamic exercises for low back pain. *Scand J Med Sci Sports* 1996;**6**:82–7.
19. **Carpenter DM**, Nelson BW. Low back strengthening for the prevention and treatment of low back pain. *Med Sci Sports Exerc* 1999;**31**:18–24.
20. **Helmhout PH**, Harts CC, Staal JB, *et al.* Comparison of a high-intensity and a low-intensity lumbar extensor training program as minimal intervention treatment in low back pain: a randomized trial. *Eur Spine J* 2004;**13**:537–47.
21. **Petersen T**, Larsen K, Jacobsen S. One-year follow-up comparison of the effectiveness of McKenzie treatment and strengthening training for patients with chronic low back pain: outcome and prognostic factors. *Spine* 2007;**32**:2948–56.
22. **Helmhout PH**, Harts CC, Viechtbauer W, *et al.* Isolated lumbar extensor strengthening versus regular physical therapy in an army working population with nonacute low back pain: a randomized controlled trial. *Arch Phys Med Rehabil* 2008;**89**:1675–85.
23. **Sertopyraz F**, Eyigor S, Karapolat H, *et al.* Comparison of isokinetic exercise versus standard exercise training in patients with chronic low back pain: a randomized controlled study. *Clin Rehabil* 2009;**23**:238–47.
24. **Limke JC**, Rainville J, Peña E, *et al.* Randomized trial comparing the effects of one set vs two sets of resistance exercises for outpatients with chronic low back pain and leg pain. *Eur J Phys Rehabil Med* 2008;**44**:399–405.
25. **Schlumberger A**, Stec J, Schmidtleicher D. Single- vs. multiple-set strength training in women. *J Strength Cond Res* 2001;**15**:284–9.
26. **Rhea MR**, Alvar BA, Burkett LN, *et al.* A meta-analysis to determine the dose response for strength development. *Med Sci Sports Exerc* 2003;**35**:456–64.
27. **Peterson MD**, Rhea MR, Alvar BA. Maximizing strength development in athletes: a meta-analysis to determine the dose-response relationship. *J Strength Cond Res* 2004;**18**:377–82.
28. **Rhea MR**. Synthesizing strength and conditioning research: the meta-analysis. *J Strength Cond Res* 2004;**18**:921–3.
29. **Siff MC**, Verkoshansky Y. *Supertraining*. 6th edition. Denver, CO: Supertraining Institute 2003.
30. **Fleck SJ**, Kraemer WJ. *Designing Resistance Training Programs*. 3rd edition. Champaign, IL: Human Kinetics 2004.
31. **Baechle TR**, Earle RW. *Essentials of Strength Training and Conditioning*. 3rd edition. Champaign, IL: Human Kinetics 2008.
32. American College of Sports Medicine Position Stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 2009;**41**:687–708.
33. **Kell RT**, Asmundson GJ. A comparison of two forms of periodized exercise rehabilitation programs in the management of chronic nonspecific low-back pain. *J Strength Cond Res* 2009;**23**:513–23.
34. **Jackson JK**, Shepherd TR, Kell RT. The influence of periodized resistance training on recreationally active males with chronic nonspecific low back pain. *J Strength Cond Res* 2011;**25**:242–51.
35. **Alfredson H**, Pietilä T, Jonsson P, *et al.* Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med* 1998;**26**:360–6.
36. **Mafi N**, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 2001;**9**:42–7.
37. **Roos EM**, Engström M, Lagerquist A, *et al.* Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy – a randomized trial with 1-year follow-up. *Scand J Med Sci Sports* 2004;**14**:286–95.
38. **Langberg H**, Ellingsgaard H, Madsen T, *et al.* Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports* 2007;**17**:61–6.
39. **Maffulli N**, Walley G, Sayana MK, *et al.* Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 2008;**30**:1677–84.
40. **Gärdin A**, Movin T, Svensson L, *et al.* The long-term clinical and MRI results following eccentric calf muscle training in chronic Achilles tendinosis. *Skeletal Radiol* 2010;**39**:435–42.
41. **Silbernagel KG**, Thomeé R, Thomeé P, *et al.* Eccentric overload training for patients with chronic Achilles tendon pain – a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sports* 2001;**11**:197–206.
42. **Kingma JJ**, de Knikker R, Wittink HM, *et al.* Eccentric overload training in patients with chronic Achilles tendinopathy: a systematic review. *Br J Sports Med* 2007;**41**:e3.
43. **Woodley BL**, Newsham-West RJ, Baxter GD. Chronic tendinopathy: effectiveness of eccentric exercise. *Br J Sports Med* 2007;**41**:188–98.
44. **Meyer A**, Tumilty S, Baxter GD. Eccentric exercise protocols for chronic non-interstitial Achilles tendinopathy: how much is enough? *Scand J Med Sci Sports* 2009;**19**:609–15.
45. **Allison GT**, Purdam C. Eccentric loading for Achilles tendinopathy – strengthening or stretching? *Br J Sports Med* 2009;**43**:276–9.
46. **Purdam CR**, Jonsson P, Alfredson H, *et al.* A pilot study of the eccentric decline squat in the management of painful chronic patellar tendinopathy. *Br J Sports Med* 2004;**38**:395–7.
47. **Young MA**, Cook JL, Purdam CR, *et al.* Eccentric decline squat protocol offers superior results at 12 months compared with traditional eccentric protocol for patellar tendinopathy in volleyball players. *Br J Sports Med* 2005;**39**:102–5.
48. **Frohm A**, Saartok T, Halvorsen K, *et al.* Eccentric treatment for patellar tendinopathy: a prospective randomised short-term pilot study of two rehabilitation protocols. *Br J Sports Med* 2007;**41**:e7.
49. **Visnes H**, Hoksrud A, Cook J, *et al.* No effect of eccentric training on jumper's knee in volleyball players during the competitive season: a randomized clinical trial. *Clin J Sport Med* 2005;**15**:227–34.
50. **Jonsson P**, Alfredson H. Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: a prospective randomised study. *Br J Sports Med* 2005;**39**:847–50.
51. **Bahr R**, Fossan B, Løken S, *et al.* Surgical treatment compared with eccentric training for patellar tendinopathy (Jumper's Knee). A randomized, controlled trial. *J Bone Joint Surg Am* 2006;**88**:1689–98.
52. **Kongsgaard M**, Kovanen V, Aagaard P, *et al.* Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sports* 2009;**19**:790–802.
53. **Kongsgaard M**, Qvortrup K, Larsen J, *et al.* Fibril morphology and tendon mechanical properties in patellar tendinopathy: effects of heavy slow resistance training. *Am J Sports Med* 2010;**38**:749–56.
54. **O'Reilly SC**, Jones A, Muir KR, *et al.* Quadriceps weakness in knee osteoarthritis: the effect on pain and disability. *Ann Rheum Dis* 1998;**57**:588–94.
55. **Stueltjens MP**, Dekker J, van Baar ME, *et al.* Muscle strength, pain and disability in patients with osteoarthritis. *Clin Rehabil* 2001;**15**:331–41.
56. **Ettlinger WH Jr**, Burns R, Messier SP, *et al.* A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997;**277**:25–31.
57. **Gür H**, Cakin N, Akova B, *et al.* Concentric versus combined concentric-eccentric isokinetic training: effects on functional capacity and symptoms in patients with osteoarthritis of the knee. *Arch Phys Med Rehabil* 2002;**83**:308–16.
58. **Topp R**, Woolley S, Hornyak J 3rd, *et al.* The effect of dynamic versus isometric resistance training on pain and functioning among adults with osteoarthritis of the knee. *Arch Phys Med Rehabil* 2002;**83**:1187–95.
59. **Lange AK**, Vanwanseele B, Fiatarone Singh MA. Strength training for treatment of osteoarthritis of the knee: a systematic review. *Arthritis Rheum* 2008;**59**:1488–94.
60. **Schilke JM**, Johnson GO, Housh TJ, *et al.* Effects of muscle-strength training on the functional status of patients with osteoarthritis of the knee joint. *Nurs Res* 1996;**45**:68–72.
61. **Mikesky AE**, Mazzuca SA, Brandt KD, *et al.* Effects of strength training on the incidence and progression of knee osteoarthritis. *Arthritis Rheum* 2006;**55**:690–9.
62. **King LC**, Birmingham TB, Kean CO, *et al.* Resistance training for medial compartment knee osteoarthritis and malalignment. *Med Sci Sports Exerc* 2008;**40**:1376–84.
63. **Jan MH**, Lin JJ, Liao JJ, *et al.* Investigation of clinical effects of high- and low-resistance training for patients with knee osteoarthritis: a randomized controlled trial. *Phys Ther* 2008;**88**:427–36.

64. **Lucchinetti E**, Adams CS, Horton WE Jr, *et al*. Cartilage viability after repetitive loading: a preliminary report. *Osteoarthr Cartil* 2002;**10**:71–81.
65. **Andersen LL**, Magnusson SP, Nielsen M, *et al*. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Phys Ther* 2006;**86**:683–97.
66. **Lange AK**, Vanwanseele B, Foroughi N, *et al*. Resistive Exercise for Arthritic Cartilage Health (REACH): a randomized double-blind, sham-exercise controlled trial. *BMC Geriatr* 2009;**9**:1.
67. **Barber-Westin SD**, Noyes FR, Heckmann TP, *et al*. The effect of exercise and rehabilitation on anterior-posterior knee displacements after anterior cruciate ligament autograft reconstruction. *Am J Sports Med* 1999;**27**:84–93.
68. **Kvist J**. Rehabilitation following anterior cruciate ligament injury: current recommendations for sports participation. *Sports Med* 2004;**34**:269–80.
69. **Myer GD**, Paterno MV, Ford KR, *et al*. Neuromuscular training techniques to target deficits before return to sport after anterior cruciate ligament reconstruction. *J Strength Cond Res* 2008;**22**:987–1014.
70. **McHugh MP**, Tyler GW, Gleim GW, *et al*. Preoperative indicators of motion loss and weakness following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 1998;**27**:407–11.
71. **Urbach D**, Nebelung W, Becker R, *et al*. Effects of reconstruction of the anterior cruciate ligament on voluntary activation of quadriceps femoris a prospective twitch interpolation study. *J Bone Joint Surg Br* 2001;**83**:1104–10.
72. **McHugh MP**, Tyler TF, Nicholas SJ, *et al*. Electromyographic analysis of quadriceps fatigue after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2001;**31**:25–32.
73. **Drechsler WI**, Cramp MC, Scott OM. Changes in muscle strength and EMG median frequency after anterior cruciate ligament reconstruction. *Eur J Appl Physiol* 2006;**98**:613–23.
74. **Ageberg E**, Roos HP, Silbernagel KG, *et al*. Knee extension and flexion muscle power after anterior cruciate ligament reconstruction with patellar tendon graft or hamstring tendons graft: a cross-sectional comparison 3 years post surgery. *Knee Surg Sports Traumatol Arthrosc* 2009;**17**:162–9.
75. **Risberg MA**, Holm I, Tjomsland O, *et al*. Prospective study of changes in impairments and disabilities after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 1999;**29**:400–12.
76. **Palmieri-Smith RM**, Thomas AC, Wojtyls EM. Maximizing quadriceps strength after ACL reconstruction. *Clin Sports Med* 2008;**27**:405–24, vii–ix.
77. **Mikkelsen C**, Werner S, Eriksson E. Closed kinetic chain alone compared to combined open and closed kinetic chain exercises for quadriceps strengthening after anterior cruciate ligament reconstruction with respect to return to sports: a prospective matched follow-up study. *Knee Surg Sports Traumatol Arthrosc* 2000;**8**:337–42.
78. **Morrissey MC**, Drechsler WI, Morrissey D, *et al*. Effects of distally fixated versus nondistally fixated leg extensor resistance training on knee pain in the early period after anterior cruciate ligament reconstruction. *Phys Ther* 2002;**82**:35–43.
79. **Holm L**, Esmarck B, Mizuno M, *et al*. The effect of protein and carbohydrate supplementation on strength training outcome of rehabilitation in ACL patients. *J Orthop Res* 2006;**24**:2114–23.
80. **Liu-Ambrose T**, Taunton JE, MacIntyre D, *et al*. The effects of proprioceptive or strength training on the neuromuscular function of the ACL reconstructed knee: a randomized clinical trial. *Scand J Med Sci Sports* 2003;**13**:115–23.
81. **Risberg MA**, Holm I, Myklebust G, *et al*. Neuromuscular training versus strength training during first 6 months after anterior cruciate ligament reconstruction: a randomized clinical trial. *Phys Ther* 2007;**87**:737–50.
82. **Meyers MC**, Sterling JC, Marley RR. Efficacy of stairclimber versus cycle ergometry in postoperative anterior cruciate ligament rehabilitation. *Clin J Sport Med* 2002;**12**:85–94.
83. **Gerber JP**, Marcus RL, Dibble LE, *et al*. Safety, feasibility, and efficacy of negative work exercise via eccentric muscle activity following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2007;**37**:10–18.
84. **Gerber JP**, Marcus RL, Dibble LE, *et al*. Effects of early progressive eccentric exercise on muscle structure after anterior cruciate ligament reconstruction. *J Bone Joint Surg Am* 2007;**89**:559–70.
85. **Gerber JP**, Marcus RL, Dibble LE, *et al*. Effects of early progressive eccentric exercise on muscle size and function after anterior cruciate ligament reconstruction: a 1-year follow-up study of a randomized clinical trial. *Phys Ther* 2009;**89**:51–9.
86. **Culliford DJ**, Maskell J, Beard DJ, *et al*. Temporal trends in hip and knee replacement in the United Kingdom: 1991 to 2006. *J Bone Joint Surg Br* 2010;**92**:130–5.
87. **Sicard-Rosenbaum L**, Light KE, Behrman AL. Gait, lower extremity strength, and self-assessed mobility after hip arthroplasty. *J Gerontol A Biol Sci Med Sci* 2002;**57**:M47–51.
88. **Trudelle-Jackson E**, Emerson R, Smith S. Outcomes of total hip arthroplasty: a study of patients one year postsurgery. *J Orthop Sports Phys Ther* 2002;**32**:260–7.
89. **Hauer K**, Specht N, Schuler M, *et al*. Intensive physical training in geriatric patients after severe falls and hip surgery. *Age Ageing* 2002;**31**:49–57.
90. **Suetta C**, Magnusson SP, Rosted A, *et al*. Resistance training in the early postoperative phase reduces hospitalization and leads to muscle hypertrophy in elderly hip surgery patients – a controlled, randomized study. *J Am Geriatr Soc* 2004;**52**:2016–22.
91. **Suetta C**, Aagaard P, Rosted A, *et al*. Training-induced changes in muscle CSA, muscle strength, EMG, and rate of force development in elderly subjects after long-term unilateral disuse. *J Appl Physiol* 2004;**97**:1954–61.
92. **Suetta C**, Andersen JL, Dalgas U, *et al*. Resistance training induces qualitative changes in muscle morphology, muscle architecture, and muscle function in elderly postoperative patients. *J Appl Physiol* 2008;**105**:180–6.
93. **Husby VS**, Helgerud J, Bjørgen S, *et al*. Early maximal strength training is an efficient treatment for patients operated with total hip arthroplasty. *Arch Phys Med Rehabil* 2009;**90**:1658–67.
94. **Husby VS**, Helgerud J, Bjørgen S, *et al*. Early postoperative maximal strength training improves work efficiency 6–12 months after osteoarthritis-induced total hip arthroplasty in patients younger than 60 years. *Am J Phys Med Rehabil* 2010;**89**:304–14.
95. **Hvid L**, Aagaard P, Justesen L, *et al*. Effects of aging on muscle mechanical function and muscle fiber morphology during short-term immobilization and subsequent retraining. *J Appl Physiol* 2010;**109**:1628–34.
96. **Monteiro AG**, Aoki MS, Evangelista AL, *et al*. Nonlinear periodization maximizes strength gains in split resistance training routines. *J Strength Cond Res* 2009;**23**:1321–6.
97. **Rhea MR**, Alderman BL. A meta-analysis of periodized versus nonperiodized strength and power training programs. *Res Q Exerc Sport* 2004;**75**:413–22.